

SUGAR BEET (*Beta vulgaris*)  
Rhizomania; *Beet necrotic yellow vein virus*  
Storage rot; *Athelia*-like sp., *Botrytis cinerea*,  
and *Penicillium* spp.

C. A. Strausbaugh, USDA-ARS NWISRL, 3793 N. 3600 E.,  
Kimberly, ID 83341

### Commercial sugar beet cultivars evaluated for rhizomania resistance and storability in Idaho, 2016.

Twenty-two commercial sugar beet (*Beta vulgaris* L.) cultivars and two rhizomania susceptible check cultivars were evaluated in a sprinkler-irrigated sugar beet field near Kimberly, ID where barley was grown in 2015. The trial was conducted in a field that contained Portneuf silt loam soil and relied on natural infection for rhizomania development. The field was fall plowed and in the spring, fertilized (90 lb N and 110 lb P<sub>2</sub>O<sub>5</sub>/A) and roller harrowed on 4 Apr 16. The plots were planted on 18 Apr to a density of 142,560 seeds/A, and thinned to 47,520 plants/A on 27 May. Plots were four rows (22-in. row spacing) and 24-ft long. The experimental design was a randomized complete block design with six replications. The crop was managed according to standard cultural practices in southern Idaho. The plots were rated for rhizomania foliar symptom (percentage of plants with yellow, stunted, upright leaves) development on 19 Aug. The plants were mechanically topped and the center two rows were dug with a mechanical harvester on 4-5 Oct. At harvest, the roots were evaluated for rhizomania using a scale of 0 to 9 (0 = healthy and 9 = dead) in a continuous manner (Plant Dis. 93:632-638). The percent sucrose at harvest was established based on two eight-root samples from each plot. The samples were submitted to The Amalgamated Sugar Co. Tare Lab (determined percent sucrose, conductivity, nitrates, and tare). At harvest, eight roots per plot were also placed in a mesh onion bag, weighed, and placed in an indoor commercial sugar beet storage facility in Paul, ID on 6 Oct set to hold 34°F. On 13 Feb 17, roots were retrieved after 130 days in storage and evaluated for surface root rot (% of root surface area), weight, and percent sucrose using gas chromatography (Plant Dis. 92:581-587). Only samples from the same plot were compared when establishing percent reduction in sucrose at harvest versus storage. Data were analyzed using the general linear models procedure (Proc GLM-SAS 9.4), and Fisher's protected least significant difference (LSD;  $\alpha = 0.05$ ) was used for mean comparisons. The foliar data were arc sine square root transformed prior to analysis, but the non-transformed means are presented in the table.

Root rots and other disease problems other than rhizomania were not evident in the plot area. There were significant differences among cultivars for all variables. Rhizomania was uniform based on foliar symptoms (100%) in the susceptible checks, BTS4D20 and C-209. Most cultivars exhibited rhizomania resistance based on foliar symptoms, except for SX026 which had 50% susceptible plants. SX026 and HM126457 were the commercial cultivars that ranked the highest for rhizomania root rating, but they were still significantly better than the susceptible checks. The greatest average root yield for any cultivar was 45.08 t/A, which was similar to Idaho's average of 41.4 t/A (USDA-National Ag. Stat. Service). The primary fungal growth was an *Athelia*-like basidiomycete (Mycologia 104:70-78), but *Botrytis cinerea* Pers. and *Penicillium* spp. (*P. expansum* Link and *P. cellarum* C.A. Strausb. & Dugan) were also frequently present. After 130 days in storage, surface root rot ranged from 7 to 51%, weight loss ranged from 7.5 to 12.3%, sucrose losses ranged from 23 to 62%, and estimated recoverable sucrose (ERS) after storage ranged from 2,901 to 10,114 lb/A. Given these response ranges, selecting cultivars for rhizomania resistance and combining this resistance with storability will lead to considerable economic benefit for the sugar beet industry. If cultivars with the greatest sucrose reduction are considered for production in the future, they should only be directly processed (early harvest cultivars) and not stored based on data for root rot and sucrose losses.

Cultivar <sup>z</sup>	Rhizomania rating <sup>y</sup>		Surface root rot (%) <sup>x</sup>	Weight reduction (%) <sup>w</sup>	Root yield (t/A)	ERS at harvest (lb/A) <sup>v</sup>	Sucrose reduction (%) <sup>u</sup>	ERS after storage (lb/A)
	Foliar (%)	Root						
B-48	1 cd	2.2 de	12 de	8.5 d-g	41.17 de	13,378 a-c	24 ij	10,114 a
B-46	0 e	2.2 d-f	10 de	9.6 c-g	41.11 de	13,126 a-d	26 h-j	9,644 ab
B-7	3 c	2.3 d	10 e	7.8 fg	40.15 e	12,478 c-f	23 j	9,591 a-c
SX021	0 de	2.2 d-g	13 de	7.9 fg	41.67 de	13,199 a-c	28 g-j	9,450 a-d
C-47	1 c-e	2.0 gh	15 de	7.9 fg	44.71 ab	13,554 ab	31 e-j	9,437 a-d
SV025	0 de	2.2 d-h	10 de	8.3 e-g	42.02 c-e	13,552 ab	30 f-j	9,426 a-d
SV012	2 c	2.2 d-g	15 de	9.6 c-g	41.84 de	12,902 a-f	28 g-j	9,300 a-e
C-34	1 c-e	2.2 de	13 de	8.8 d-g	40.98 de	12,655 b-f	29 g-j	8,961 a-f
SV016	0 e	2.1 d-h	10 de	8.4 d-g	42.67 a-e	12,964 a-f	35 d-j	8,551 a-g
SV026	0 de	2.1 d-h	16 c-e	9.9 b-f	43.15 a-d	13,691 a	37 c-i	8,544 a-g
B-110	1 cd	2.2 d-f	14 de	8.2 fg	40.42 e	12,794 a-f	35 d-j	8,336 a-h
C-29	0 e	2.1 d-h	13 de	10.5 a-e	42.74 a-e	13,065 a-e	39 c-h	8,061 b-h
C-36	0 e	2.0 gh	7 e	8.9 d-g	44.51 a-c	13,353 a-c	40 b-h	8,058 b-h
C-204	0 de	2.0 h	22 c-e	8.3 e-g	45.08 a	13,602 a	43 b-f	7,808 c-h
HM 12SYN003	0 de	2.2 d-g	22 c-e	9.7 c-g	41.67 de	12,129 f	37 c-i	7,693 d-h
HM 103425	2 c	2.2 d-h	22 c-e	10.6 a-d	42.30 b-e	12,225 d-f	39 c-h	7,510 e-h
HM 118711	1 cd	2.2 d-g	17 c-e	10.5 a-e	42.55 a-e	12,603 c-f	41 b-g	7,507 e-h
C-48	0 e	2.0 f-h	9 e	7.5 g	44.60 a-c	13,292 a-c	44 b-e	7,365 f-h
HM 126457	1 c	2.6 c	45 ab	9.4 d-g	34.12 f	10,706 g	34 d-j	6,948 gh
HM 125891	0 de	2.1 e-h	51 a	11.8 a-c	43.30 a-d	12,201 ef	45 b-d	6,653 h
SX026	50 b	2.6 c	17 c-e	8.8 d-g	33.98 f	10,212 g	36 d-j	6,528 hi
C-39	1 cd	2.2 d-h	45 ab	12.0 ab	42.38 b-e	12,526 c-f	62 a	4,778 ij
<b>BTS4D20</b>	100 a	3.8 b	25 cd	11.7 a-c	29.47 g	8,029 h	52 ab	3,822 jk
<b>C-209</b>	100 a	4.1 a	31 bc	12.3 a	21.03 h	5,744 i	50 a-c	2,901 k
<i>P</i> > <i>F</i> <sup>t</sup>	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001
LSD ( $\alpha = 0.05$ )	Trans	0.2	15	2.2	2.65	901	13	1,811

<sup>z</sup> For more information on coded cultivars, contact the following companies: B = Betaseed Inc., C = ACH Seeds Inc., HM = Hillehog, SV = SESVanderHave, and SX = Seedex. Rhizomania susceptible check cultivars were BTS4D20 and C-209 (Bold).

<sup>y</sup> Foliar = percentage of foliage in plot with rhizomania symptoms on 19 Aug. Root = roots were evaluated for rhizomania using a scale of 0 to 9 (0 = healthy, 9 = dead; Plant Dis. 93:632-638) in a continuous manner at harvest.

<sup>x</sup> Surface root rot = percentage of root surface area discolored in storage.

<sup>w</sup> Weight reduction = difference in weight from harvest to the end of storage.

<sup>v</sup> ERS = estimated recoverable sucrose was calculated as extraction  $\times$  0.01  $\times$  gross sucrose and extraction = 250 + [1,255.2  $\times$  (conductivity - 15,000)  $\times$  (percent sucrose - 6,185)] / (percent sucrose  $\times$  [98.66 - (7.845  $\times$  conductivity)]).

<sup>u</sup> Sucrose reduction (%) = (1 - ((% Sucrose<sub>storage sample</sub> - 1.395)  $\times$  Weight<sub>storage sample</sub>) / (% Sucrose<sub>harvest sample</sub>  $\times$  Weight<sub>harvest sample</sub>))  $\times$  100.

<sup>t</sup> *P* > *F* was the probability associated with the *F* value. Within each variable, means followed by the same letter did not differ significantly based on Fisher's protected least significant difference (LSD;  $\alpha = 0.05$ ). Trans = the foliar data were arc sine square root transformed prior to analysis, but the non-transformed means are presented in the table.