

SUGAR BEET (*Beta vulgaris*)
Rhizomania; *Beet necrotic yellow vein virus*
Storage rot; *Athelia*-like sp., *Botrytis cinerea*,
and *Penicillium* spp.

C. A. Strausbaugh, USDA-ARS NWISRL, 3793 N. 3600 E.,
Kimberly, ID 83341

Commercial sugar beet cultivars evaluated for rhizomania resistance and storability in Idaho, 2019.

Thirty-seven commercial sugar beet (*Beta vulgaris* L.) cultivars and two rhizomania susceptible check cultivars were evaluated in a sprinkler-irrigated sugar beet field near Kimberly, ID where barley was grown in 2018. The trial was conducted in a field that contained Portneuf silt loam soil and relied on natural infection for rhizomania development. The field was plowed and fertilized (90 lb N and 110 lb P₂O₅/A) and roller harrowed on 11 Apr 19. The plots were planted on 23 Apr to a density of 142,560 seeds/A, and thinned to 47,520 plants/A on 5 Jun. Plots were four rows (22-in. between-row spacing) and 24-ft long. The experimental design was a randomized complete block design with six replications. The crop was managed according to standard cultural practices in southern Idaho. The plots were rated for rhizomania foliar symptom (percentage of plants with yellow, stunted, upright leaves) development on 20 Aug. The plants were mechanically topped and the center two rows were dug with a mechanical harvester on 7-8 Oct. At harvest, the roots were evaluated for rhizomania symptoms using a scale of 0 to 9 (0 = healthy and 9 = dead) in a continuous manner (Plant Dis. 93:632-638). The percent sucrose at harvest was established based on two eight-root samples from each plot. The samples were submitted to The Amalgamated Sugar Co. Tare Lab (determined percent sucrose, conductivity, nitrates, and tare). At harvest, eight roots per plot were also placed in a mesh onion bag, weighed, and placed in an indoor commercial sugar beet storage facility in Paul, ID on 9 Oct set to hold 34°F. On 2 Mar 20, roots were retrieved after 145 days in storage and evaluated for surface root rot (% of root surface area), weight, and percent sucrose using high performance liquid chromatography (Plant Dis. 92:581-587). Only samples from the same plot were compared when establishing percent reduction in sucrose at harvest versus storage. Data were analyzed using the general linear models procedure (Proc GLM-SAS 9.4), and Fisher's protected least significant difference (LSD; $\alpha = 0.05$) was used for mean comparisons. The foliar data were arc sine square root transformed and root rating data were rank transformed prior to analysis, but the non-transformed means are presented in the table.

Root rots and other disease problems other than rhizomania were not evident in the plot area. There were significant differences among cultivars for all variables. Rhizomania was uniform based on foliar symptoms (100%) in the susceptible checks, BTS4D20 and C-209. Most cultivars exhibited rhizomania resistance based on foliar symptoms, except for five cultivars (B48, C47, HM103425, HM118711, and MA715) which had 31 to 80% susceptible plants. Cultivars HM103425 and HM118711 also had root ratings that were not significantly different from the susceptible checks. The highest average root yield for any cultivar was 41.8 t/A, which was similar to Idaho's average of 39.0 t/A (USDA-National Ag. Stat. Service). The primary fungal growth in storage was an *Athelia*-like basidiomycete (Mycologia 104:70-78), but *Botrytis cinerea* Pers. and *Penicillium* spp. (*P. expansum* Link and *P. cellarum* C.A. Strausb. & Dugan) were also frequently present. After 145 days in storage, surface root rot ranged from 10 to 68%, weight loss ranged from 14 to 25%, sucrose reduction ranged from 22 to 72%, and estimated recoverable sucrose (ERS) after storage ranged from 2,442 to 9,769 lb/A. Given these response ranges, selecting cultivars for rhizomania resistance and combining this resistance with storability will lead to considerable economic benefit for the sugar beet industry. If cultivars with the highest sucrose reduction are considered for production in the future, they should only be directly processed (early harvest cultivars) and not stored based on data for root rot and sucrose reduction.

Cultivar ^z	Rhizomania rating ^y		Surface root rot (%) ^x	Weight reduction (%) ^w	Root yield (t/A)	ERS at harvest (lb/A) ^v	Sucrose reduction (%) ^u	ERS after storage (lb/A)
	Foliar (%)	Root						
B71	1 i	2.4 f-n	10 p	14.2 kl	38.96 c-l	12,591 b-h	22 r	9,769 a
SV032	1 i	2.2 n-r	13 op	21.1 bc	40.85 a-e	13,350 ab	30 pq	9,357 ab
B79	0 i	2.3 l-r	13 n-p	16.7 e-l	40.63 a-f	13,479 a	31 m-q	9,246 ab
SX036	1 hi	2.2 m-r	14 m-p	14.4 j-l	41.43 a-c	13,377 ab	32 k-q	9,062 a-c
B66	0 i	2.3 k-r	15 m-p	18.4 c-i	40.22 a-g	13,021 a-e	31 n-q	8,987 a-d
C54	0 i	2.4 f-o	30 f-i	15.7 h-l	39.31 b-j	12,422 c-i	28 qr	8,933 a-d
B74	2 g-i	2.2 p-r	38 d-f	16.6 e-l	41.22 a-d	13,347 ab	33 j-q	8,883 a-e
C49	0 i	2.2 p-r	16 l-p	18.1 c-i	41.43 a-c	13,230 a-c	34 i-q	8,735 b-f
C57	4 f-i	2.4 g-o	46 c-e	13.7 l	38.59 e-l	12,372 d-i	30 o-q	8,599 b-g
SV026	2 g-i	2.1 r	28 f-k	17.2 d-l	41.82 a	13,158 a-d	37 f-n	8,241 c-h
C58	0 i	2.4 i-q	18 j-p	15.8 h-l	39.26 b-k	12,415 c-i	34 h-q	8,195 c-h
SV039	2 g-i	2.6 e-j	24 g-n	17.9 c-j	37.74 g-m	12,122 f-j	33 k-q	8,178 c-i
B65	1 i	2.2 o-r	17 l-p	18.7 c-h	41.42 a-c	13,101 a-d	38 f-l	8,132 c-j
B69	1 hi	2.4 h-p	25 g-m	19.4 b-g	40.03 a-h	12,395 d-i	35 g-q	8,057 d-j
C48	2 g-i	2.4 g-o	30 f-i	18.7 c-h	39.24 b-k	12,224 e-j	35 g-p	7,929 e-k
B73	1 i	2.6 d-h	15 l-p	17.4 d-k	36.53 lm	11,509 j-n	32 l-q	7,870 f-k
SV028	3 f-i	2.5 f-m	17 k-p	16.3 f-l	38.40 e-l	12,555 b-h	38 f-n	7,860 f-k
SX026	3 f-i	2.6 f-k	26 g-l	15.8 h-l	38.22 f-l	12,240 e-j	36 f-p	7,830 f-k
B78	1 hi	2.5 f-k	32 f-h	15.1 i-l	38.12 g-l	12,188 f-j	37 f-o	7,694 g-l
SX021	3 g-i	2.5 f-n	24 g-n	16.0 g-l	38.51 e-l	12,233 e-j	38 f-m	7,611 h-l
C204	2 g-i	2.2 qr	34 fg	15.7 h-l	41.50 ab	12,622 b-g	40 e-i	7,536 h-m
SX030	1 g-i	2.4 i-q	20 i-p	17.2 d-l	38.86 d-l	12,375 d-i	40 e-j	7,453 h-n
SV033	3 g-i	2.5 f-l	22 h-o	13.7 l	38.74 d-l	12,588 b-h	42 d-f	7,365 h-n
B48	48 d	2.6 d-i	13 op	17.4 d-k	37.44 i-m	11,868 g-k	38 f-k	7,316 h-o
SV025	2 g-i	2.5 f-k	17 l-p	17.3 d-k	38.25 f-l	12,244 e-j	41 d-g	7,224 i-o
SX034	3 f-i	2.4 j-q	29 f-j	19.4 b-g	39.59 a-i	12,507 c-h	43 d-f	7,178 j-o
HIL9919NT	3 f-i	2.5 f-k	24 g-n	17.7 c-k	38.76 d-l	11,828 g-k	41 d-h	7,006 k-o
HIL9916	8 f-h	2.6 d-i	30 f-i	20.1 b-e	37.78 g-m	11,783 h-l	43 d-f	6,756 l-o
HM12SYN003	2 g-i	2.6 e-j	39 d-f	19.4 b-g	37.83 g-m	10,770 n	38 f-k	6,613 m-p
C47	69 c	2.6 b-f	32 f-h	22.6 ab	37.20 i-m	11,001 l-n	40 d-i	6,565 n-q
C56	1 i	2.6 c-g	30 f-i	19.0 c-h	37.31 i-m	11,830 g-k	46 de	6,361 o-q
MA715	31 e	2.8 a-e	34 e-g	16.2 f-l	35.60 mn	10,873 mn	47 d	5,734 pq
HM126457	9 fg	3.7 a	64 a	17.8 c-j	28.14 p	8,853 op	36 f-p	5,627 q
HM125891	2 g-i	3.2 a-c	67 a	20.6 b-d	32.24 o	9,072 o	54 c	4,221 r
BTS4D20	100 a	3.8 a	48 b-d	19.6 b-f	29.42 p	8,145 p	57 c	3,569 rs
C39	10 f	2.6 c-g	52 bc	18.1 c-i	37.57 h-m	11,280 k-n	69 ab	3,532 rs
HM103425	80 b	3.0 a-d	58 ab	17.8 c-j	33.37 no	9,612 o	64 b	3,455 rs
HM118711	69 c	3.1 ab	68 a	17.9 c-j	32.56 o	9,558 o	72 a	2,730 st
C209	100 a	5.2 a	24 g-n	25.3 a	20.18 q	5,475 q	56 c	2,442 t
<i>P</i> > <i>F</i> ^t	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001
LSD ($\alpha = 0.05$)	Trans	Trans	11	3.5	2.53	832	7	968

^z For more information on coded cultivars, contact the following companies: B and BTS = Betaseed Inc., C = ACH Seeds Inc., HIL and HM = Hillshög, MA = Maribo, SV = SESVanderHave, and SX = Seedex. Rhizomania susceptible check cultivars were BTS4D20 and C209.

^y Foliar = percentage of foliage in plot with rhizomania symptoms on 20 Aug. Root = roots were evaluated for rhizomania using a scale of 0 to 9 (0 = healthy, 9 = dead; Plant Dis. 93:632-638) in a continuous manner at harvest.

^x Surface root rot = percentage of root surface area discolored in storage.

^w Weight reduction = difference in weight from harvest to the end of storage.

^v ERS = estimated recoverable sucrose was calculated as extraction x 0.01 x gross sucrose and extraction = 250 + [1255.2 x (conductivity - 15000) x (percent sucrose - 6185)] / (percent sucrose x [98.66 - (7.845 x conductivity)]).

^u Sucrose reduction (%) = (1 - (((% Sucrose_{storage sample} - 1.395) x Weight_{storage sample}) / (% Sucrose_{harvest sample} x Weight_{harvest sample}))) x 100.

^t *P* > *F* was the probability associated with the *F* value. Within each variable, means followed by the same letter did not differ significantly based on Fisher's protected least significant difference (LSD; $\alpha = 0.05$). Trans = the foliar data were arc sine square root transformed and the root rating data were rank transformed prior to analysis, but the non-transformed means are presented in the table.