

## Absorption of Tricarballic Acid from the Rumen of Sheep and Cattle Fed Forages Containing *trans*-Aconitic Acid

James B. Russell\*

Agricultural Research Service, United States Department of Agriculture, Ithaca, New York, USA

and Henry F. Mayland

Agricultural Research Service, United States Department of Agriculture, Kimberly, Idaho, USA

(Received 3 December 1986; revised version received 20 January 1987; accepted 29 January 1987)

### ABSTRACT

*Some forages accumulate high concentrations (<5% of dry matter) of trans-aconitate, and this acid has been implicated in Mg chelation and the occurrence of grass tetany in ruminants. In vitro experiments have indicated that rumen microorganisms convert trans-aconitate to tricarballic acid. The feeding studies described here were conducted to demonstrate absorption of tricarballic acid by ruminant animals fed diets similar to those producing grass tetany. When sheep were switched from a diet containing alfalfa (lucerne) (Medicago sativa L.) hay (no detectable trans-aconitate) to wheat (Triticum aestivum L.) and rye (Secale cereale L.) forage containing 1.52 and 1.37% trans-aconitate, respectively, there was a rapid increase in blood plasma tricarballic acid. Trans-aconitate was not detected in the plasma. At 16 h after feeding, plasma tricarballic acid concentrations were  $0.58 \pm 0.08$  and  $0.48 \pm 0.21$  mM in sheep fed the wheat and rye forage, respectively. Tricarballic acid concentrations remained relatively constant for the remaining 60 h of the experiment. Cattle were fed rye forage one week later, and the concentration of trans-aconitate in the forage had dropped to 0.83% of the dry matter. Once again there was a rapid*

\*Address for correspondence: J. B. Russell, 321 Morrison Hall, Cornell University, Ithaca, NY 14853, USA.

appearance of tricarballylate in plasma, but the maximum concentration was  $0.31 \pm 0.05$  mM ( $t=27$  h). When the cattle were removed from the rye forage, there was a linear decline in tricarballylate and none was detected 24 h later. The studies indicated that *trans*-aconitate is converted to tricarballylate in the rumen and that tricarballylate rather than *trans*-aconitate is absorbed.

**Key words:** Tricarballylate, aconitate, magnesium, grass tetany.

## 1 INTRODUCTION

Magnesium deficiency of ruminants has been recognized since the 1930s and the symptoms have been identified by a variety of names including grass tetany, lactation tetany, and grass staggers. The disease is most often observed in lactating cattle and sheep grazing succulent, cool-season grass species.<sup>1-4</sup> The availability of forage Mg is influenced by a number of factors including low concentrations of nonstructural carbohydrates and high concentrations of K, N, higher fatty acids, and other organic acids.<sup>5</sup> The finding of large amounts of *trans*-aconitate in forages known to induce grass tetany has led several researchers<sup>6-8</sup> to suggest that *trans*-aconitate might chelate Mg and reduce the availability of dietary Mg. Oral administration of *trans*-aconitate or citrate in conjunction with KCl induced grass tetany symptoms that were corrected by Ca and Mg therapy.<sup>9</sup> The symptoms did not occur when the animals were given orally either acid independent of the KCl. In another study, *trans*-aconitate disappeared rapidly from the sheep rumen, but little *trans*-aconitate was detected in either blood or urine.<sup>10</sup>

*In vitro* experiments indicated that *trans*-aconitate was rapidly fermented by rumen microorganisms.<sup>11</sup> Some of the *trans*-aconitate was converted to acetate but as much as 50% was converted to tricarballylate (Fig. 1). Tricarballylate was a fermentation product of *trans*-aconitate but not of other organic acids *in vitro*. Tricarballylate was not metabolized further by rumen microorganisms, and it appeared that tricarballylate rather than *trans*-aconitate would remain in the rumen for a long enough time to affect Mg absorption.<sup>11</sup> Sheep given gelatin capsules filled with *trans*-aconitate absorbed significant amounts of tricarballylate, not aconitate, into blood.<sup>12</sup> *In vitro* studies with purified enzymes and isolated hepatocytes indicated that tricarballylate was an inhibitor of the enzyme aconitase and could inhibit acetate oxidation in the citric acid cycle.<sup>12</sup>

The formation of tricarballylate was previously verified with animals fed timothy hay (*Phleum pratense* L.).<sup>11,12</sup> Since this forage did not contain detectable amounts of *trans*-aconitate, reagent-grade *trans*-aconitate was provided as a supplement. The following experiments were performed in the western United States where *trans*-aconitate-accumulating plants are the predominant range forages.<sup>7</sup> Feeding trials were initiated during the early spring when lush grasses usually contain the highest concentration of *trans*-aconitate.




Fig. 1. Structure and likely pathway of aconitate conversion to tricarballylate. Taken from reference 11 with permission.

## 2 EXPERIMENTAL

### 2.1 Feeding trials

Six nonlactating, nonpregnant, 6-year-old white-face ewes (56 kg) and two lactating 2-year-old Suffolk ewes (61 kg), each with twin lambs (24 kg), were fed alfalfa (*Medicago sativa* L.) at 0500 and 1600 for 4 days prior to diet change. The ewes were randomly assigned to one of two treatment groups with one lactating ewe in each group. One group received freshly harvested wheat (*Triticum aestivum* L.) forage (Table 1) which was in the preboot stage of growth (23% dry matter), and the other group was given rye (*Secale cereale* L.) in the early preboot stage (19% dry matter). The forage was fed at 0500 and 1600 for the duration of the experiment, and water was provided *ad libitum*. A second experiment was conducted with four lactating beef cows weighing approximately 410 kg. These animals were also fed alfalfa hay for 4 days before the change in diet. During the experimental period (1.5 days), cows were fed (0500 and 1600) freshly harvested rye forage.

**TABLE 1**  
Concentrations of *trans*-Aconitate, Sugars and Minerals in Forages Fed to Sheep and Cattle

Animal	Forage	Forage composition							K	Ash <sup>a</sup> alkalinity (meq kg <sup>-1</sup> )	Dry matter <sup>b</sup> intake (kg)	Aconitate <sup>b</sup> intake (mm)
		Aconitate	Glucose	Fructose	Sucrose	Ca	Mg	K				
		(% dry matter)										
Ewes	Alfalfa hay	0.00	2.30	0.70	0.80	1.29	0.32	2.00	0.56	0	<i>ad lib.</i>	—
	Rye	1.37	1.40	1.28	11.19	0.31	0.13	2.86	2.80	626	4.1	330
	Wheat	1.52	1.24	1.10	14.21	0.31	0.13	2.92	2.85	629	4.6	410
Cows	Alfalfa hay	0.00	1.10	0.60	1.10	1.27	0.34	2.38	0.66	0	<i>ad lib.</i>	—
	Rye	0.83	1.40	1.02	11.95	0.28	0.13	2.89	3.00	640	12.5	610

<sup>a</sup>Residual ash corrected for nitrate loss which is a good estimate of total organic acids.<sup>20</sup>

<sup>b</sup>Intake for the experimental feeding period of 76 and 32 h for ewes and cows, respectively.




Fig. 2. Effect of forage ( $\Delta$ , wheat, or  $\blacktriangle$ , rye) on the concentration of tricarballylate in the blood of sheep (a) and cattle (b). The animals were switched from alfalfa to rye or wheat at time 0 ( $\blacktriangledown$ ). The cattle were switched back to lucerne at 24 h ( $\blacktriangleleft$ ).

## 2.2 Analyses

Blood was drawn by jugular puncture into heparinized tubes (see Fig. 2 for interval). The blood was centrifuged, and the plasma was frozen and preserved during transfer from Kimberly, Idaho, to Ithaca, New York, for analysis. Representative forage samples were dried at 60°C for 48 h to calculate dry matter content, and the samples were ground to pass through a 1-mm sieve.

Forage samples were extracted with 0.026 M  $H_2SO_4$  (24 h at 5°C) and centrifuged (10 000 $\times g$ , 15 min, 0°C) to remove insoluble forage residue. The forage extract and blood samples were deproteinized with  $HClO_4$  (0.5 M, final concentration) and the  $ClO_4^-$  was removed with an excess of  $K_2CO_3$ . Organic acids and sugars were measured by high-pressure liquid chromatography using a Beckman model 334 liquid chromatograph that was equipped with a model 156 refractive index detector and a Bio Rad HPX-87H organic acid column.<sup>13</sup> The sample size was 50  $\mu l$ , the eluent was 0.026 M  $H_2SO_4$ , and the column temperature was 50°C. Magnesium, Ca and K were measured by atomic absorption spectrophotometry.<sup>14</sup>

## 3 RESULTS AND DISCUSSION

During the background period the sheep were fed alfalfa hay that did not contain any detectable aconitate (Table 1). At time zero the sheep were given either wheat or rye forage, and the intake of *trans*-aconitate was 410 and 330 mM, respectively, over the next 72 h. Soon after feeding there was a rapid increase in

blood tricarballoylate (Fig. 2a), and aconitate was not detected in any of the blood samples. Within 2 h tricarballoylate concentrations in plasma had reached 0.2 mM, and by 16 h the concentration was 0.58 mM with the wheat and 0.48 mM for the rye forage. During the next 60 h, tricarballoylate ranged from 0.44 to 0.60 mM. Tricarballoylate absorption did not affect ( $P < 0.05$ ) plasma concentrations of Mg or Ca, and tetany was not observed. During the background period, the concentrations of plasma Mg and Ca ( $n = 16$ ) were  $24.8 \pm 0.4$  and  $113.4 \pm 1.4$  mg liter<sup>-1</sup>, respectively. After the change in diet, the values ( $n = 56$ ) were  $24.8 \pm 0.4$  and  $115 \pm 1.4$  mg liter, respectively. In this study the intake of *trans*-aconitate was probably lower than amounts consumed by many animals that have developed grass tetany. Grasses sometimes contain more than 5% *trans*-aconitate.<sup>6</sup>

The cattle study was conducted one week after the sheep study, and the aconitate content of the rye forage had declined to 0.83% (Table 1). When the cattle were switched from alfalfa to rye, plasma concentrations of tricarballoylate increased rapidly (Fig. 2b). There was only 60% as much aconitate in the rye during this trial as in the previous trial, and the maximum concentration of tricarballoylate in plasma was half that observed in the sheep trial (Fig. 2). When the cattle were switched back to alfalfa hay, blood concentrations of tricarballoylate declined rapidly. While being fed the rye diet, the cows ingested 610 mg *trans*-aconitate. Plasma Mg and Ca were  $24.4 \pm 0.9$  and  $117.6 \pm 2.7$  mg liter<sup>-1</sup>, respectively, during the background period and did not change significantly after tricarballoylate was absorbed ( $23.0 \pm 0.09$  and  $122 \pm 1.2$ , respectively).

Previous studies indicated that *trans*-aconitate was rapidly metabolized to tricarballoylate.<sup>11,12</sup> The almost immediate appearance of tricarballoylate in the plasma of sheep and cattle in the study reported here is consistent with the earlier observations. The forages used in this study were not particularly high in *trans*-aconitate; however, plasma concentrations of tricarballoylate were greater than 0.5 mM in the sheep study (Fig. 2a). Since the plasma concentration seemed to be dose dependent (sheep versus cattle trial), it is conceivable that plasma concentrations could be greater in ruminants grazing forage with even more *trans*-aconitate.

The wheat and rye forages contained high concentrations of sucrose in addition to *trans*-aconitate (Table 1), and the sugar may have enhanced tricarballoylate formation. When rumen bacteria capable of reducing aconitate to tricarballoylate were isolated, *Selenomonas ruminantium* was the predominant species.<sup>15</sup> *S. ruminantium* has an extremely high affinity for sucrose ( $K_m = 4 \mu M$ ), and its numbers increase greatly when sugars are present in the rumen.<sup>16,17</sup> Bacteria were not enumerated in this study, but, on the basis of sucrose content in the wheat and rye forage, the numbers of aconitate-reducing bacteria should have increased during the experimental period.

When cattle were returned to the alfalfa diet, tricarballoylate concentrations in the blood declined (Fig. 2b). Disappearance could have been attributed to either metabolism or excretion, but the latter seemed more likely. Tricarballoylate is an effective metabolic inhibitor and it is not rapidly metabolized by mammalian enzymes.<sup>12</sup> Renal clearance rates<sup>18</sup> could account for the disappearance of tricarballoylate.

#### 4 CONCLUSIONS

These experiments provide evidence that sheep and cattle fed forages containing *trans*-aconitate converted the *trans*-aconitate to tricarballic acid, another acid capable of chelating magnesium (stability constant of 115).<sup>19</sup> Since tricarballic acid rather than *trans*-aconitate was absorbed, tricarballic acid could be a significant factor in the etiology of 'grass tetany'. Further experiments should be conducted to ascertain blood concentrations of tricarballic acid in tetanic animals, the ability of tricarballic acid to inhibit the citric acid cycle, and the capacity of tricarballic acid to increase urinary losses of magnesium.

#### REFERENCES

1. Fontenot, J. P. Animal nutrition aspects of grass tetany. In: *Grass Tetany* (Rendig, V. V.; Grunes, D. L., Eds), American Society for Agronomy, Madison, WI., USA, 1979, 50-62.
2. Littledike, E. T.; Cox, P. S. Clinical, mineral, and endocrine interrelationships in hypomagnemic tetany. In: *Grass Tetany* (Rendig, V. V.; Grunes, D. L., Eds), American Society for Agronomy, Madison, WI., USA, 1979, 1-50.
3. Mayland, H. F.; Grunes, D. L. Soil-climate-plant relationships in the etiology of grass tetany. In: *Grass Tetany* (Rendig, V. V.; Grunes, D. L., Eds), American Society for Agronomy, Madison, Wis., USA, 1979, 123-175.
4. McDonald, P.; Edwards, R. A.; Greenhalgh, J. F. D. Minerals. In: *Animal Nutrition*, Hefner Press, Edinburgh, 1973, 98-100.
5. Mayland, H. F.; Grunes, D. L. Soil-climate-plant relationships in the etiology of grass tetany. In: *Grass Tetany* (Rendig, V. V.; Grunes, D. L., Eds), American Society for Agronomy, Madison, WI., 1979, 123-175.
6. Burau, R.; Stout, P. R. *Trans*-aconitic acid in range grasses in early spring. *Science* 1965, **150**, 766-767.
7. Stout, P. R.; Brownell, J.; Burau, R. G. Occurrences of *trans*-aconitate in range forage species. *Agron. J.* 1967, **59**, 21-24.
8. Scotto, K. C.; Bohman, V. R.; Lesperance, A. L. Effects of oral potassium and sodium chloride on plasma composition of cattle: a grass tetany related study. *J. Anim. Sci.* 1971, **32**, 354-358.
9. Bohman, V. R.; Lesperance, A. L.; Harding, G. D.; Grunes, D. L. Induction of experimental tetany in cattle. *J. Anim. Sci.* 1969, **29**, 99-102.
10. Kennedy, G. S. *Trans*-aconitate utilization by sheep. *Aust. J. Biol. Sci.* 1968, **21**, 529-538.
11. Russell, J. B.; Van Soest, P. J. *In vitro* ruminal fermentation of organic acids common in forage. *Appl. Environ. Microbiol.* 1984, **47**, 155-159.
12. Russell, J. B.; Forsberg, N. Production of tricarballic acid by rumen microorganisms and its potential toxicity in ruminant tissue metabolism. *Br. J. Nutr.* 1986, **56**, 153-162.
13. Erlich, G. G.; Goerlitz, D. F.; Bowell, J. H.; Eisen, G. V.; Godsy, E. M. Liquid chromatographic procedure for fermentation product analysis in the identification of anaerobic bacteria. *Appl. Environ. Microbiol.* 1980, **40**, 608-612.
14. Perkin-Elmer. *Analytical Methods for Atomic Absorption Spectrophotometry*, Perkin-Elmer, Norwalk, CT, 1982.
15. Russell, J. B. Enrichment and isolation of rumen bacteria that reduce *trans*-aconitic acid to tricarballic acid. *Appl. Environ. Microbiol.* 1985, **49**, 120-126.
16. Bryant, M. P. The characteristics of strains of *Selenomonas* isolated from bovine rumen contents. *J. Bacteriol.* 1956, **72**, 162-167.

17. Russell, J. B.; Baldwin, R. L. Comparison of substrate affinities among several rumen bacteria: a possible determinant of rumen bacterial competition. *Appl. Environ. Microbiol.* 1979, **37**, 531-536.
18. Guyton, A. C. *Textbook of Medical Physiology*, 4th edn, W. B. Saunders Co., Philadelphia, PA, 1971, 406-414.
19. Martell, A. E.; Smith, R. M. *Critical Stability Constants*, vol. 3, Plenum Press, New York, 1977.
20. van Tuil, H. D.; Lampe, J. E. M.; Dijkshoorn, W. The possibility of relating the ash alkalinity to organic-salt content. *Jaarboek I.B.S., Wageningen, The Netherlands*, 1964, **250**, 157-160.

Library of Congress  
Washington, D.C. 20540

U.S. Government Printing Office  
Washington, D.C. 20540

1980  
101-101